

HEPATITIS C

General Information

What is hepatitis?

“Hepatitis” means inflammation of the liver. The liver is a vital organ that processes nutrients, filters the blood, and fights infections. When the liver is inflamed or damaged, its function can be affected.

Heavy alcohol use, toxins, some medications, and certain medical conditions can cause hepatitis. However, hepatitis is most often caused by a virus. In the United States, the most common types of viral hepatitis are Hepatitis A, Hepatitis B, and Hepatitis C.



Most people who get infected with the Hepatitis C virus develop a chronic, or lifelong, infection.

What is Hepatitis C?

Hepatitis C is an infection of the liver that results from the Hepatitis C virus. **Acute** Hepatitis C refers to the first several months after someone is infected. Acute infection can range in severity from a very mild illness with few or no symptoms to a serious condition requiring hospitalization. For reasons that are not known, about 20% of people are able to clear, or get rid of, the virus without treatment in the first 6 months.

Unfortunately, most people who get infected are not able to clear the Hepatitis C virus and develop a chronic, or lifelong, infection. Over time, **chronic** Hepatitis C can cause serious health problems including liver disease, liver failure, and even liver cancer.

How is Hepatitis C spread?

Hepatitis C is usually spread when blood from a person infected with the Hepatitis C virus enters the body of someone who is not infected. Today, most people become infected with Hepatitis C by sharing needles, syringes, or any other equipment to inject drugs. Before widespread screening of the blood supply in 1992, Hepatitis C was also spread through blood transfusions and organ transplants. While uncommon, poor infection control has resulted in outbreaks in healthcare settings.

While rare, sexual transmission of Hepatitis C is possible. Having a sexually transmitted disease or HIV, sex with multiple partners, or rough sex appears to increase a person’s risk for Hepatitis C. Hepatitis C can also be spread when getting tattoos and body piercings in unlicensed facilities, informal settings, or with non-sterile instruments. Also, approximately 6% of infants born to infected mothers will get Hepatitis C. Still, some people don’t know how or when they got infected.

What are the symptoms of Hepatitis C?

Many people with Hepatitis C do not have symptoms and do not know they are infected. If symptoms occur, they can include: fever, feeling tired, not wanting to eat, upset stomach, throwing up, dark urine, grey-colored stool, joint pain, and yellow skin and eyes.

When do symptoms occur?

If symptoms occur with acute infection, they can appear anytime from 2 weeks to 6 months after infection. If symptoms occur with chronic Hepatitis C, they can take decades to develop. When symptoms appear with chronic Hepatitis C, they often are a sign of advanced liver disease.

Continued on next page



How would you know if you have Hepatitis C?

The only way to know if you have Hepatitis C is to get tested. Doctors use a blood test, called a Hepatitis C Antibody Test, which looks for antibodies to the Hepatitis C virus. Antibodies are chemicals released into the bloodstream when someone gets infected. Antibodies remain in the bloodstream, even if the person clears the virus.

A positive or reactive Hepatitis C Antibody Test means that a person has been infected with the Hepatitis C virus at some point in time. However, a positive antibody test **does not** necessarily mean a person still has Hepatitis C. An additional test called a RNA test is needed to determine if a person is currently infected with Hepatitis C.

Who should get tested for Hepatitis C?

Testing for Hepatitis C is recommended for certain groups, including people who:

- Were born from 1945 – 1965
- Received donated blood or organs before 1992
- Have ever injected drugs, even if it was just once or many years ago
- Have certain medical conditions, such as chronic liver disease and HIV or AIDS
- Have abnormal liver tests or liver disease
- Have been exposed to blood from a person who has Hepatitis C
- Are on hemodialysis
- Are born to a mother with Hepatitis C

Can Hepatitis C be treated?

Yes. However, treatment depends on many different factors, so it is important to see a doctor experienced in treating Hepatitis C. New and improved treatments are available that can cure Hepatitis C for many people.



Testing is the only way to know if you have Hepatitis C.

How can Hepatitis C be prevented?

Although there is currently no vaccine to prevent Hepatitis C, there are ways to reduce the risk of becoming infected with the Hepatitis C virus.

- Avoid sharing or reusing needles, syringes or any other equipment to prepare and inject drugs, steroids, hormones, or other substances.
- Do not use personal items that may have come into contact with an infected person's blood, even in amounts too small to see, such as razors, nail clippers, toothbrushes, or glucose monitors.
- Do not get tattoos or body piercings from an unlicensed facility or in an informal setting.

For more information

Talk to your health professional, call your health department, or visit www.cdc.gov/hepatitis.